

## Health Care Reform Preventive Services Coding Guide

The Patient Protection and Affordable Care Act (PPACA) and the Health Care and Education Reconciliation Act of 2010 (HCERA) has designated the services listed below as preventive benefits and available with no cost-sharing when provided by an in-network provider for members of non-grandfathered health plans. In addition to the services listed below, your patient may have additional preventive care benefits covered under their health plan that may or may not be covered at 100%. Your patient should check their benefit booklet for details on these additional preventive care benefits. The following tables provide a quick reference guide for submitting claims for preventive services with a "well-person" diagnosis code as the primary (first) diagnosis on the claim. This information is intended as a reference tool for your convenience and is not a guarantee of payment. **This guide is subject to change based on new or revised laws and/or regulations, additional guidance and/or PALIC-PR medical policy.**

**IMPORTANT INFORMATION:** Services must be billed with the appropriate diagnosis, at the line level of the claim (Block 24E), pursuant to industry standard coding guidelines. Preventive or screening services are intended for those who currently exhibit no signs or symptoms of disease. Services otherwise deemed preventive that are received in an inpatient setting, an emergency room, or that include additional procedures or diagnostic services may be subject to copayment, deductible and coinsurance. Submitting screening service codes (CPT, HCPCS, ICD-9 or ICD-10) when signs or symptoms are present constitutes inappropriate coding which could result in recoupment of monies paid to the provider for those services. Additionally, these services are subject to certain limitations depending on medical necessity and other reasonable medical management techniques. If you have questions, please contact the **PALIC-PR at (787) 620-1414**.

<b>Grade A and B Recommendations of U.S. Preventive Services Task Force (USPSTF) currently effective unless otherwise noted</b>	<b>CPT or HCPCS</b>	<b>ICD-10 Diagnosis</b>	<b>Comments</b>
<b>Screening for Abdominal Aortic Aneurysm</b> (one time screening for abdominal aortic aneurysm by ultrasonography in men ages 65-75 who have ever smoked)	76706	Z13.6	
<b>Screening and counseling to reduce alcohol misuse</b> The USPSTF recommends that clinicians screen adults aged 18 years or older for alcohol misuse and provide persons engaged in risky or hazardous drinking with brief behavioral counseling interventions to reduce alcohol misuse.	99408, 99409, G0442 or G0443, G0396, G0397 <b>Effective 1/1/19:G2011</b>	Z13.89, F10.10, F10.120, F10.129	These codes are to be used in the absence of a wellness visit.
<b>Aspirin to prevent cardiovascular disease and colorectal cancer in men and women</b> (Low-dose aspirin use for the primary prevention of cardiovascular disease (CVD) and colorectal cancer (CRC) in adults aged 50 to 59 years who have a 10% or greater 10-year CVD risk, are not at increased risk for bleeding, have a life expectancy of at least 10 years, and are willing to take low-dose aspirin daily for at least 10 years.)	Not applicable, administered through Pharmacy	Not applicable, administered through Pharmacy	OTC Aspirin (81 mg) is dispensed to member with a physician order with no cost-sharing.
<b>Screening for bacteriuria</b> (screening for asymptomatic bacteriuria with <u>urine culture</u> for pregnant women at 12 - 16 weeks' gestation or at the first prenatal visit, if later)	87081, 87084, 87086 or 87088	Z34.00, Z34.01, Z34.02, Z34.03, Z34.80, Z34.81, Z34.82, Z34.83, Z34.90, Z34.91, Z34.92, Z34.93, Z33.1, O09.00, O09.01, O09.02, O09.03, O09.10, O09.11, O09.12, O09.13, O09.291, O09.292, O09.293, O09.299, O09.40, O09.41, O09.42, O09.43, O09.211, O09.212, O09.213, O09.219, O09.30, O09.31, O09.32, O09.33, O09.511, O09.512, O09.513, O09.519, O09.521, O09.522, O09.523, O09.529, O09.611, O09.612, O09.613, O09.619, O09.621, O09.622, O09.623, O09.629, O09.811, O09.812, O09.813, O09.819, O09.821, O09.822, O09.823, O09.829, O36.80X0, O36.80X1, O36.80X2, O36.80X3, O36.80X4, O36.80X5, O36.80X9, O09.891, O09.892, O09.893, O09.899, O09.70, O09.71, O09.72, O09.73, O09.90, O09.91, O09.92, O09.93, O09.A0, O09.A1, O09.A2, O09.A3	Dipstick or other urinalysis included in global maternity visits. Only urine culture part of preventive services mandate. <b>NOTE: Z33.1 is not an acceptable principle diagnosis code.</b>
<b>Screening for high blood pressure</b> (Recommends screening for high blood pressure in adults aged 18 years or older. The USPSTF recommends obtaining measurements outside of the clinical setting for diagnostic confirmation before starting treatment)	99385-99387 or 99395-99397 <b>For ambulatory blood pressure monitoring use</b> 93784, 93786, 93788, 93790 <b>For home blood pressure monitor use</b> A4670	Z13.6 or other wellness exam diagnosis code For coverage of ABPM or HBPM diagnosis code R03.0 is required	Part of wellness office visit. <b>Effective 1/1/2017</b> , ambulatory blood pressure monitoring (ABPM) or home blood pressure monitors (HBPM) for confirmation of high blood pressure is covered at 100%. Home blood pressure monitors must be purchased from an in-network durable medical equipment (DME) vendor.
<b>Counseling related to BRCA screening</b> (Recommends that primary care providers screen women who have family members with breast, ovarian, tubal, or peritoneal cancer with 1 of several screening tools designed to identify a family history that may be associated with an increased risk for potentially harmful mutations in breast cancer susceptibility genes (BRCA 1 or BRCA 2). Women with positive screening results should receive generic counseling and, if indicated after counseling, BRCA testing.)	Referrals for counseling included in wellness visit codes 99385-99387, 99395-99397 Counseling services use 96040, 99401, 99402, 99403, 99404, or S0265 Genetic testing use <del>81211</del> , 81212, <del>81213</del> , <del>81214</del> , 81215, 81216, 81217 or 81162 <b>Effective 1/1/19:</b> 81163, 81164, 81165, 81166, 81167	Z80.3, Z80.41, Z85.07, Z85.3, Z85.43, Z85.44, Z85.46, or Z71.83	Services for BRCA 1/BRCA 2 testing will be provided with no cost sharing to appropriate groups when the medical policy criteria are met. <a href="#">See PALIC-PR Medical Policy: Genetic Testing for Breast and Ovarian Cancer</a> 99401-99404 are to be used in the absence of a wellness visit.

## Health Care Reform Preventive Services Coding Guide

<b>Grade A and B Recommendations of U.S. Preventive Services Task Force (USPSTF) currently effective unless otherwise noted</b>	<b>CPT or HCPCS</b>	<b>ICD-10 Diagnosis</b>	<b>Comments</b>
<b>Screening for breast cancer [mammography]</b> (for women aged 40 or over every 1-2 yrs , with or without clinical breast examination)	77067, 77063	Z12.31, Z12.39	If the patient has had an abnormal mammogram in the past, subsequent routine mammograms may be coded as diagnostic: 77061, 77062, 77065, 77066, G0279 with a diagnosis reflective of the abnormality.
<b>Chemoprevention of breast cancer</b> (Recommends that clinicians engage in shared, informed decision making with women who are at increased risk for breast cancer about medications to reduce their risk. For women who are at increased risk for breast cancer and at low risk for adverse medication effects, clinicians should offer to prescribe risk-reducing medications, such as tamoxifen or raloxifene)	99401-99402 or is included in wellness visit 99385-99387, 99395-99397	Z15.01, Z80.3, D24.1, D24.2, D24.9, N60.81, N60.82, N60.89	This recommendation applies to asymptomatic women aged 35 years or older without a prior diagnosis of breast cancer, ductal carcinoma in situ, or lobular carcinoma in situ. Generic risk-reducing medications are dispensed to member with a physician order with no cost-sharing. For members who have swallowing problems or may have an intolerance to generic products, brand products may also be made available by completing the <b>Copay Waiver Form</b> and faxing it to the number on the bottom of the document. 99401-99404 are to be used in the absence of a wellness visit.  NOTE: Z15.01 is not an acceptable principle diagnosis
<b>Interventions to support breast feeding</b> (interventions during pregnancy and after birth to promote and support breastfeeding)	99401-99403 or part of other office visit	O30.93, Z34.03, Z34.83, Z34.93, Z39.1	99401-99403 are to be used in the absence of a wellness visit.
<b>Screening for cervical cancer</b> (in women ages 21 to 65 yrs with cytology (pap smear) every 3 yrs or, for women ages 30-65 yrs who want to lengthen the screening interval, screening with a combination of cytology and human papillomavirus (HPV) testing every 5 yrs)	88141, 88142, 88143, 88147, 88148, 88150, 88152, 88153, 88164, 88165, 88166, 88167, 88174, 88175, 87623, 87624, 87625, G0101, G0123, G0124, G0141, G0143, G0144, G0145, G0147, G0148, P3000, P3001	Z00.00, Z00.01, Z01.411, Z01.419, Z01.42, Z12.4, Z11.51	This recommendation does not apply to women who have received a diagnosis of a high-grade precancerous cervical lesion or cervical cancer, women with in utero exposure to diethylstilbestrol, or women who are immunocompromised (such as those who are HIV positive).
<b>Screening for chlamydial infection in non-pregnant and pregnant women</b> (for all sexually active non-pregnant and pregnant young women ages 24 and younger and for older, non-pregnant and pregnant women who are at increased risk)	87270, 87320, 87490, 87491 and 87810	Z34.00, Z34.01, Z34.02, Z34.03, Z34.80, Z34.81, Z34.82, Z34.83, Z34.90, Z34.91, Z34.92, Z34.93, Z33.1, O09.00, O09.01, O09.02, O09.03, O09.10, O09.11, O09.12, O09.13, O09.291, O09.292, O09.293, O09.299, O09.40, O09.41, O09.42, O09.43, O09.211, O09.212, O09.213, O09.219, O09.30, O09.31, O09.32, O09.33, O09.511, O09.512, O09.513, O09.519, O09.521, O09.522, O09.523, O09.529, O09.611, O09.612, O09.613, O09.619, O09.621, O09.622, O09.623, O09.629, O09.811, O09.812, O09.813, O09.819, O09.821, O09.822, O09.823, O09.829, O36.80X0, O36.80X1, O36.80X2, O36.80X3, O36.80X4, O36.80X5, O36.80X9, O09.891, O09.892, O09.893, O09.899, O09.70, O09.71, O09.72, O09.73, O09.90, O09.91, O09.92, O09.93, Z11.8, O09.A0, O09.A1, O09.A2, O09.A3, Z11.3	<b>NOTE: Z33.1 is not an acceptable principle diagnosis code.</b>
<b>Screening for cholesterol abnormalities:</b> <b>*men 35 and older</b> (for lipid disorders) <b>*men younger than 35</b> (for ages 20-35 for lipid disorders if they are at increased risk for coronary heart disease) <b>*women 20 and older</b> (for lipid disorders if they are at increased risk for coronary heart disease)  <b>See also: Statin Use for the Primary Prevention of Cardiovascular Disease in Adults</b>	80061, 82465 or 83718	Z13.220	Do not bill the panel lab code in addition to separate tests included in the panel.

## Health Care Reform Preventive Services Coding Guide

<p><b>Grade A and B Recommendations of U.S. Preventive Services Task Force (USPSTF) currently effective unless otherwise noted</b></p>	<p><b>CPT or HCPCS</b></p>	<p><b>ICD-10 Diagnosis</b></p>	<p><b>Comments</b></p>
<p><b>Prevention of dental caries</b> (recommends that primary care clinicians prescribe oral fluoride supplementation starting at age 6 months up to age 16 for children whose water supply is deficient in fluoride. Also recommends that primary care clinicians apply fluoride varnish to the primary teeth of all infants and children starting at the age of primary tooth eruption.)</p>	<p>99188, 99401 or 99402 May prescribe oral fluoride during child wellness visit: 99381-99382 or 99391-99392 with appropriate wellness diagnosis</p>	<p>Z29.3</p>	<p>Fluoride varnish will not be covered beyond the 5th birthday. Oral fluoride supplementation will be covered from ages 6 months through 16 years.</p> <p>PALIC-PR recommends counseling/education in addition to varnish application.</p> <ul style="list-style-type: none"> <li>• For well child visit: Any counseling services received at the time of a well child visit are considered part of the preventive visit and are not reimbursed separately.</li> <li>• For sick visit: If the varnish is applied during a visit other than for wellness, providers should reference diagnosis code Z41.8 or <b>Z29.3</b> as the primary diagnosis for the line item of the varnish application service, 99188, on the claim in order to process correctly. If education/counseling on prevention and risk factor reduction of dental caries is conducted during a sick visit, providers should again reference diagnosis code Z41.8 or <b>Z29.3</b> as the primary diagnosis for the line item of the counseling service, 99401/99402, on the claim in order to process correctly.</li> </ul> <p>PALIC-PR will disallow D codes if submitted by non licensed dental providers 99401-99404 are to be used in the absence of a wellness visit.</p>
<p><b>Screening for colorectal cancer</b> (using fecal occult blood testing, sigmoidoscopy, or colonoscopy, in adults beginning at ages 50-75)</p>	<p><b>Fecal occult blood testing:</b> 81528, 82270, 82274 or G0328 <b>Sigmoidoscopy:</b> 45330, 45333, 45334, 45338, G0104, G6022, 88305 and <b>Effective 1/1/2017</b> G0500 <b>Colonoscopy:</b> 45378, 45380, 45381, 45382, 45384, 45385, 74263, G0105, G0121, G6024, 00812, 88305 and G0500 <b>Consultation or office visit prior to screening colonoscopy:</b> S0285</p>	<p><b>Screening Procedures:</b> Z12.11, Z12.12, K51.40, K62.1, K63.5, D12.0, D12.2, D12.3, D12.4, D12.5, D12.6, D12.7, D12.8, or D12.9 <b>Pathology services (88305):</b> Z12.11, Z12.12, K51.40, K62.1, K63.5, D12.0, D12.2, D12.3, D12.4, D12.5, D12.6, D12.7, D12.8, or D12.9 <b>Visit or Consultations:</b> Z01.818</p>	<p>Colorectal cancer screening mandate does not include barium enema. ***If a test is performed for screening purposes, screening service codes may be submitted even in the event of positive findings e.g., polyp found during a screening colonoscopy. It is critical to use a screening diagnosis as the primary diagnosis for the claim line of the test performed [not just in the header diagnoses]. Any abnormal findings from a screening test should be listed as a secondary diagnosis code. Failure to list screening as the primary diagnosis code will affect correct claim adjudication. When billing for a visit or consultation prior to a screening colonoscopy, S0285 should be submitted with the appropriate diagnosis code. <b>As of 9/1/2016</b>, Certain <a href="#">bowel preparation medications</a>, when medically appropriate and prescribed by a health care provider are allowed without cost sharing when dispensed at a participating pharmacy. If there is a medical reason a member cannot take a generic bowel preparation, the physician should review the clinical <a href="#">criteria</a> and if appropriate, submit fax <a href="#">form</a>. <b>As of 1/1/2017</b>, Moderate sedation (G0500) performed by the physician performing the procedure is allowed but cannot be billed on the same DOS as 00812.</p>
<p><b>Screening for depression:</b> <b>*adults</b> (screening for depression, including pregnant and postpartum women, when staff-assisted depression care supports are in place to assure accurate diagnosis, effective treatment, and follow-up) <b>*adolescents</b> (screening 12-18 yr olds for major depressive disorder when systems are in place to ensure accurate diagnosis, psychotherapy [cognitive-behavioral or interpersonal], and follow-up)</p>	<p>G0444 or preventive visit</p>	<p><b>Z13.89</b> Remove effective 9/30/18 <b>Effective 10/1/18: Z13.31</b></p>	<p>Part of any problem or preventive office visit.</p>



## Health Care Reform Preventive Services Coding Guide

<p><b>Grade A and B Recommendations of U.S. Preventive Services Task Force (USPSTF) currently effective unless otherwise noted</b></p>	<p><b>CPT or HCPCS</b></p>	<p><b>ICD-10 Diagnosis</b></p>	<p><b>Comments</b></p>
<p><b>Screening for diabetes</b> Screening for abnormal blood glucose as part of cardiovascular risk assessment. Clinicians should offer or refer patients with abnormal blood glucose to intensive behavioral counseling interventions to promote a healthful diet and physical activity. Adults aged 40 to 70 who are overweight or obese</p>	<p>82947 or 83036</p>	<p>Z13.1,</p>	
<p><b>Behavioral Counseling to promote a healthful diet and physical activity for cardiovascular disease prevention in adults with cardiovascular risk factors</b> (Recommends offering or referring adults who are overweight or obese and have additional cardiovascular disease (CVD) risk factors to intensive behavioral counseling interventions to promote a healthful diet and physical activity for CVD prevention.)</p>	<p>99402, 99403, 99404, 97802, 97803, 97804, S9452, S9470, G0270, G0271, G0108, G0109, S9140, S9141, S9455, S9460, S9465, G0446 or part of preventive visit</p>	<p>Z71.3</p>	<p>99401-99404 are to be used in the absence of a wellness visit.</p>
<p><b>Fall Prevention</b> (Exercise interventions to prevent falls in community-dwelling adults 65 years or older who are at increased risk for falls.)</p>	<p>97001-97002, 97110, 97112, 97116, 97530, G0151, G0157, G0159, S9131 and S9476.</p>	<p>Z91.81</p>	<p>Vitamin D is dispensed to member with a physician order with no cost-sharing.</p>
<p><b>Supplementation with folic acid</b> (for all women planning or capable of pregnancy to take a daily supplement containing 0.4 - 0.8 mg [400-600 mcg] of folic acid)</p>	<p>Not applicable, administered through Pharmacy</p>	<p>Not applicable, administered through Pharmacy</p>	<p>OTC folic acid supplements are dispensed to member with a physician order with no cost-sharing.</p>
<p><b>Screening for Gestational Diabetes Mellitus</b> (Recommends screening for gestational diabetes mellitus (GDM) in asymptomatic pregnant women after 24 weeks of gestation.)</p>	<p>82950</p>	<p>Z34.00, Z34.01, Z34.02, Z34.03, Z34.80, Z34.81, Z34.82, Z34.83, Z34.90, Z34.91, Z34.92, Z34.93, Z33.1, O09.00, O09.01, O09.02, O09.03, O09.10, O09.11, O09.12, O09.13, O09.291, O09.292, O09.293, O09.299, O09.40, O09.41, O09.42, O09.43, O09.211, O09.212, O09.213, O09.219, O09.30, O09.31, O09.32, O09.33, O09.511, O09.512, O09.513, O09.519, O09.521 O09.522, O09.523, O09.529, O09.611, O09.612, O09.613, O09.619, O09.621, O09.622, O09.623, O09.629, O09.811, O09.812, O09.813, O09.819, O09.821, O09.822, O09.823, O09.829, O36.80X0, O36.80X1, O36.80X2, O36.80X3, O36.80X4, O36.80X5, O36.80X9, O09.891, O09.892, O09.893, O09.899, O09.70, O09.71, O09.72, O09.73, O09.90, O09.91, O09.92, O09.93 O09.A0, O09.A1, O09.A2, O09.A3</p>	<p>In pregnant women between 24 and 28 weeks of gestation and at the first prenatal visit for pregnant women identified to be at high risk for diabetes <b>NOTE: Z33.1 is not an acceptable principle diagnosis code.</b></p>
<p><b>Screening for gonorrhea for women</b> (all sexually active women, including those who are pregnant if they are at increased risk for infection [if they are young or have other individual or population risk factors])</p>	<p>87590-87591 and 87850</p>	<p>Z11.59, Z11.3, Z34.00, Z34.01, Z34.02, Z34.03, Z34.80, Z34.81, Z34.82, Z34.83, Z34.90, Z34.91, Z34.92, Z34.93, Z33.1, O09.00, O09.01, O09.02, O09.03, O09.10, O09.11, O09.12, O09.13, O09.291, O09.292, O09.293, O09.299, O09.40, O09.41, O09.42, O09.43, O09.211, O09.212, O09.213, O09.219, O09.30, O09.31, O09.32, O09.33, O09.511, O09.512, O09.513, O09.519, O09.521 O09.522, O09.523, O09.529, O09.611, O09.612, O09.613, O09.619, O09.621, O09.622, O09.623, O09.629, O09.811, O09.812, O09.813, O09.819, O09.821, O09.822, O09.823, O09.829, O36.80X0, O36.80X1, O36.80X2, O36.80X3, O36.80X4, O36.80X5, O36.80X9, O09.891, O09.892, O09.893, O09.899, O09.70, O09.71, O09.72, O09.73, O09.90, O09.91, O09.92, O09.93, Z11.8 O09.A0, O09.A1, O09.A2, O09.A3</p>	<p><b>NOTE: Z33.1 is not an acceptable principle diagnosis code.</b></p>
<p><b>Prophylactic medication for gonorrhea: newborns</b> (ocular topical medication for all newborns against gonococcal ophthalmia neonatorum)</p>	<p>Not applicable, administered through Facility</p>	<p>Not applicable, administered through Facility</p>	<p>This medication is generally administered to newborn at birth facility.</p>
<p><b>Screening for hearing loss</b> (in all newborn infants)</p>	<p>92558, 99381, 99391 or 92586</p>	<p>Z00.121, Z00.129, Z00.110, Z00.111, Z01.10, Z01.118</p>	<p>Service is typically performed in the birth facility or as part of a wellness office visit in the event of a home birth. CPT code <b>92586</b> cannot be billed by professional provider in a facility setting</p>
<p><b>Screening for hemoglobinopathies</b> (for sickle cell disease in newborns)</p>	<p>83020, 83021, 85660 and S3620</p>	<p>Z13.0</p>	<p>Service is typically performed in the birth facility or as part of a wellness office visit in the event of a home birth.</p>

## Health Care Reform Preventive Services Coding Guide

<p><b>Grade A and B Recommendations of U.S. Preventive Services Task Force (USPSTF) currently effective unless otherwise noted</b></p>	<p><b>CPT or HCPCS</b></p>	<p><b>ICD-10 Diagnosis</b></p>	<p><b>Comments</b></p>
<p><b>Screening for Hepatitis B</b> (in pregnant women at first prenatal visit and in persons at high risk for infection.)</p>	<p>80055 , 87340 or 80081</p>	<p>Z34.00, Z34.01, Z34.02, Z34.03, Z34.80, Z34.81, Z34.82, Z34.83, Z34.90, Z34.91, Z34.92, Z34.93, Z33.1, O09.00, O09.01, O09.02, O09.03, O09.10, O09.11, O09.12, O09.13, O09.291, O09.292, O09.293, O09.299, O09.40, O09.41, O09.42, O09.43, O09.211, O09.212, O09.213, O09.219, O09.30, O09.31, O09.32, O09.33, O09.511, O09.512, O09.513, O09.519, O09.521 O09.522, O09.523, O09.529, O09.611, O09.612, O09.613, O09.619, O09.621, O09.622, O09.623, O09.629, O09.811, O09.812, O09.813, O09.819, O09.821, O09.822, O09.823, O09.829, O36.80X0, O36.80X1, O36.80X2, O36.80X3, O36.80X4, O36.80X5, O36.80X9, O09.891, O09.892, O09.893, O09.899, O09.70, O09.71, O09.72, O09.73, O09.90, O09.91, O09.92, O09.93, Z11.59 O09.A0, O09.A1, O09.A2, O09.A3</p>	<p><b>NOTE: Z33.1 is not an acceptable principle diagnosis code.</b></p>
<p><b>Screening for hepatitis C virus (HCV) infection</b> (for persons at high risk for infection.) The USPSTF also recommends offering one-time screening for HCV infection to adults born between 1945 and 1965.</p>	<p>G0472</p>	<p>Z72.51, Z72.52, Z72.53, Z11.59, Z11.4, F10.10, F10.20, F10.21, F10.229, F11.10, F11.20, F11.21, F12.10, F12.20, F12.21, F13.10, F13.20, F13.21, F14.10, F14.20, F14.21, F15.10, F15.129, F15.20, F15.21, F16.10, F16.129, F16.20, F16.21, F17.200, F18.10, F19.10, F19.20, F19.21, F55.2</p>	
<p><b>Screening for HIV</b> (on all adolescents ages 15 to 65 years, younger adolescents and older adults who are at increased risk, and all pregnant women including those who present in labor who are untested and whose HIV status is unknown)</p>	<p>G0432, G0433, G0435, OR G0475</p>	<p>Z34.00, Z34.01, Z34.02, Z34.03, Z34.80, Z34.81, Z34.82, Z34.83, Z34.90, Z34.91, Z34.92, Z34.93, Z33.1, O09.00, O09.01, O09.02, O09.03, O09.10, O09.11, O09.12, O09.13, O09.291, O09.292, O09.293, O09.299, O09.40, O09.41, O09.42, O09.43, O09.211, O09.212, O09.213, O09.219, O09.30, O09.31, O09.32, O09.33, O09.511, O09.512, O09.513, O09.519, O09.521 O09.522, O09.523, O09.529, O09.611, O09.612, O09.613, O09.619, O09.621, O09.622, O09.623, O09.629, O09.811, O09.812, O09.813, O09.819, O09.821, O09.822, O09.823, O09.829, O36.80X0, O36.80X1, O36.80X2, O36.80X3, O36.80X4, O36.80X5, O36.80X9, O09.891, O09.892, O09.893, O09.899, O09.70, O09.71, O09.72, O09.73, O09.90, O09.91, O09.92, O09.93, Z11.4 O09.A0, O09.A1, O09.A2, O09.A3</p>	<p><b>NOTE: Z33.1 is not an acceptable principle diagnosis code.</b></p>
<p><b>Screening for congenital hypothyroidism</b> (in newborns)</p>	<p>99381 or 99391</p>	<p>Z13.29</p>	<p>Service is typically performed in the birth facility or as part of a wellness office visit in the event of a home birth.</p>
<p><b>Screening for iron deficiency anemia</b> (in asymptomatic pregnant women)</p>	<p>80055, 85013, 85014, 85018, 85025, 85027 or 80081</p>	<p>Z34.00, Z34.01, Z34.02, Z34.03, Z34.80, Z34.81, Z34.82, Z34.83, Z34.90, Z34.91, Z34.92, Z34.93, Z33.1, O09.00, O09.01, O09.02, O09.03, O09.10, O09.11, O09.12, O09.13, O09.291, O09.292, O09.293, O09.299, O09.40, O09.41, O09.42, O09.43, O09.211, O09.212, O09.213, O09.219, O09.30, O09.31, O09.32, O09.33, O09.511, O09.512, O09.513, O09.519, O09.521 O09.522, O09.523, O09.529, O09.611, O09.612, O09.613, O09.619, O09.621, O09.622, O09.623, O09.629, O09.811, O09.812, O09.813, O09.819, O09.821, O09.822, O09.823, O09.829, O36.80X0, O36.80X1, O36.80X2, O36.80X3, O36.80X4, O36.80X5, O36.80X9, O09.891, O09.892, O09.893, O09.899, O09.70, O09.71, O09.72, O09.73, O09.90, O09.91, O09.92, O09.93 O09.A0, O09.A1, O09.A2, O09.A3</p>	<p><b>NOTE: Z33.1 is not an acceptable principle diagnosis code.</b></p>
<p><b>Iron supplementation in children</b> (routine iron supplementation for asymptomatic children 6-12 mo of age who are at increased risk for iron deficiency anemia)</p>	<p>Not applicable, administered through Pharmacy</p>	<p>Not applicable, administered through Pharmacy</p>	<p>OTC iron supplements are dispensed to member with a physician order with no cost-sharing.</p>
<p><b>Screening for latent tuberculosis</b> (asymptomatic adults 18 years and older at increased risk for tuberculosis)</p>	<p>86480, 86481, 86580</p>	<p>Z11.1</p>	<p>Mandate effective 10/1/2017</p>
<p><b>Low-Dose Aspirin Use for the Prevention of Morbidity and Mortality From Preeclampsia</b> (Recommends the use of low-dose aspirin (81 mg/d) as preventive medication after 12 weeks of gestation in women who are at high risk for preeclampsia.)</p>	<p>Not applicable, administered through Pharmacy</p>	<p>Not applicable, administered through Pharmacy</p>	<p>OTC Aspirin (81 mg) is dispensed to member with a physician order at a participating pharmacy with no cost-sharing.</p>

## Health Care Reform Preventive Services Coding Guide

<b>Grade A and B Recommendations of U.S. Preventive Services Task Force (USPSTF) currently effective unless otherwise noted</b>	<b>CPT or HCPCS</b>	<b>ICD-10 Diagnosis</b>	<b>Comments</b>
<b>Screening for Lung Cancer</b> (Recommends annual screening for lung cancer with low-dose computed tomography (LDCT) in adults aged 55 to 80 yrs who have a 30 pack-year smoking history and currently smoke or have quite within the past 15 years. Screening should be discontinued once a person has not smoked for 15 years or develops a health problem that substantially limits life expectancy or the ability or willingness to have curative lung surgery.)	G0297	F17.210, F17.211, F17.213, F17.218, F17.219, Z72.0, and Z87.891	
<b>Screening for and Management of Obesity in Adults</b> (screen all adults; clinicians should offer or refer patients with a body mass index (BMI) of 30 kg/m <sup>2</sup> or higher to intensive, multi-component behavioral interventions.)	99385, 99386, 99387, 99395, 99396, 99397, 96150, 96151, 96152, 96153, 99401, 99402, 99403, 99404, G0447 or G0473	Z00.00, Z00.01, Z13.89, Z68.30, Z68.31, Z68.32, Z68.33, Z68.34, Z68.35, Z68.36, Z68.37, Z68.38, Z68.39, Z68.41, Z68.42, Z68.43, Z68.44, Z68.45, E66.9, E66.01, E66.09, E66.1, E66.8, Z13.220, Z13.228, Z13.29	99401-99404 are to be used in the absence of a wellness visit. <b>NOTE: Z68.30, Z68.31, Z68.32, Z68.33, Z68.34, Z68.35, Z68.36, Z68.37, Z68.38, Z68.39, Z68.41, Z68.42, Z68.43, Z68.44, Z68.45 are not acceptable principle diagnosis codes.</b>
<b>Screening and counseling for obesity: children</b> (screen children aged 6 yrs and older for obesity and offer/refer to comprehensive, intensive behavioral interventions to promote improvement in weight status)	99383-99384, 99393, 99394, 99401, 99402, 99403, 99404, G0447 or G0473	Z00.121, Z00.129, Z00.110, Z00.111, Z13.89, Z13.220, Z13.228, Z13.29	99401-99404 are to be used in the absence of a wellness visit.
<b>Screening for osteoporosis</b> (screen women aged 65 and older for osteoporosis and in younger postmenopausal women who are at increased risk of osteoporosis, as determined by a formal clinical risk assessment tool.)	77080 or 77081	Z13.820	This recommendation uses only DEXA scans for testing.
<b>Screening for PKU</b> (in newborns)	84030	Z13.228	Service is typically performed in the birth facility or as part of a wellness office visit in the event of a home birth. 84030 is to be used in the absence of a panel which includes this test.
<b>Screening for preeclampsia</b>	Performed as part of routine prenatal visit	Z34.00, Z34.01, Z34.02, Z34.03, Z34.80, Z34.81, Z34.82, Z34.83, Z34.90, Z34.91, Z34.92, Z34.93, Z33.1, O09.00, O09.01, O09.02, O09.03, O09.10, O09.11, O09.12, O09.13, O09.291, O09.292, O09.293, O09.299, O09.40, O09.41, O09.42, O09.43, O09.211, O09.212, O09.213, O09.219, O09.30, O09.31, O09.32, O09.33, O09.511, O09.512, O09.513, O09.519, O09.521, O09.522, O09.523, O09.529, O09.611, O09.612, O09.613, O09.619, O09.621, O09.622, O09.623, O09.629, O09.811, O09.812, O09.813, O09.819, O09.821, O09.822, O09.823, O09.829, O36.80X0, O36.80X1, O36.80X2, O36.80X3, O36.80X4, O36.80X5, O36.80X9, O09.891, O09.892, O09.893, O09.899, O09.70, O09.71, O09.72, O09.73, O09.90, O09.91, O09.92, O09.93, O09.A0, O09.A1, O09.A2, O09.A3	<b>Effective 5/1/18 or upon renewal</b> Service is performed as part of routine prenatal visit not separately reimbursable <b>NOTE: Z33.1 is not an acceptable principle diagnosis code.</b>
<b>Screening for Rh incompatibility: *first pregnancy visit</b> (recommends Rh (D) blood typing and antibody testing for all pregnant women during their first visit for pregnancy-related care) <b>*24-28 weeks gestation</b> (recommends repeated Rh (D) antibody testing for all unsensitized Rh (D)-negative women at 24-28 weeks gestation unless the biological father is known to be Rh (D)-negative)	86850, 80055 or 80081	Z34.00, Z34.01, Z34.02, Z34.03, Z34.80, Z34.81, Z34.82, Z34.83, Z34.90, Z34.91, Z34.92, Z34.93, Z33.1, O09.00, O09.01, O09.02, O09.03, O09.10, O09.11, O09.12, O09.13, O09.291, O09.292, O09.293, O09.299, O09.40, O09.41, O09.42, O09.43, O09.211, O09.212, O09.213, O09.219, O09.30, O09.31, O09.32, O09.33, O09.511, O09.512, O09.513, O09.519, O09.521, O09.522, O09.523, O09.529, O09.611, O09.612, O09.613, O09.619, O09.621, O09.622, O09.623, O09.629, O09.811, O09.812, O09.813, O09.819, O09.821, O09.822, O09.823, O09.829, O36.80X0, O36.80X1, O36.80X2, O36.80X3, O36.80X4, O36.80X5, O36.80X9, O09.891, O09.892, O09.893, O09.899, O09.70, O09.71, O09.72, O09.73, O09.90, O09.91, O09.92, O09.93, O09.A0, O09.A1, O09.A2, O09.A3	Initial testing is part of the obstetric panel. Use 86850 only if performed separately from the panel for repeat testing. <b>NOTE: Z33.1 is not an acceptable principle diagnosis code.</b>
<b>Counseling for Sexually Transmitted Infections</b> (recommends high-intensity behavioral counseling to prevent STIs for all active adolescents and for adults at increased risk for STIs)	G0445, 99383, 99384, 99385, 99386, 99387, 99393, 99394, 99395, 99396, 99397 or 99401, 99402, 99403, 99404	Z71.7, Z71.89, Z72.51, Z72.52, Z72.53	99401-99404 are to be used in the absence of a wellness visit.

## Health Care Reform Preventive Services Coding Guide

<b>Grade A and B Recommendations of U.S. Preventive Services Task Force (USPSTF) currently effective unless otherwise noted</b>	<b>CPT or HCPCS</b>	<b>ICD-10 Diagnosis</b>	<b>Comments</b>
<b>Behavioral Counseling to Prevent Skin Cancer</b> (Counseling young adults, adolescents, children, and parents of young children about minimizing exposure to UV radiation for persons aged 6 months to 24 years with fair skin types to reduce their risk of skin cancer)	99383, 99384, 99385, 99393, 99394, 99395	Z00.121, Z00.129, Z00.110, Z00.111, Z00.00, Z00.01	Considered part of wellness office visit.
<b>Screening for syphilis:</b> * <b>non-pregnant persons</b> (screen persons at increased risk for syphilis infection) * <b>pregnant women</b> (screen all for syphilis infection)	80055, 86592, 86780 or 80081	Z34.00, Z34.01, Z34.02, Z34.03, Z34.80, Z34.81, Z34.82, Z34.83, Z34.90, Z34.91, Z34.92, Z34.93, Z33.1, O09.00, O09.01, O09.02, O09.03, O09.10, O09.11, O09.12, O09.13, O09.291, O09.292, O09.293, O09.299, O09.40, O09.41, O09.42, O09.43, O09.211, O09.212, O09.213, O09.219, O09.30, O09.31, O09.32, O09.33, O09.511, O09.512, O09.513, O09.519, O09.521, O09.522, O09.523, O09.529, O09.611, O09.612, O09.613, O09.619, O09.621, O09.622, O09.623, O09.629, O09.811, O09.812, O09.813, O09.819, O09.821, O09.822, O09.823, O09.829, O36.80X0, O36.80X1, O36.80X2, O36.80X3, O36.80X4, O36.80X5, O36.80X9, O09.891, O09.892, O09.893, O09.899, O09.70, O09.71, O09.72, O09.73, O09.90, O09.91, O09.92, O09.93, Z72.51, Z72.52, Z72.53, Z11.3 O09.A0, O09.A1, O09.A2, O09.A3	<b>NOTE: Z33.1 is not an acceptable principle diagnosis code.</b>
<b>Statin Use for the Primary Prevention of Cardiovascular Disease in Adults</b> (Initiate use of low- to moderate-dose statins in adults aged 40 to 75 years without a history of CVD who have 1 or more CVD risk factors (dyslipidemia, diabetes, hypertension, or smoking) and a calculated 10-year CVD event risk of 10% or greater)	Not applicable, administered through Pharmacy	Not applicable, administered through Pharmacy	Medications covered at 100%: Lovastatin (20 or 40 mg) or Pravastatin (10, 20 40 or 80 mg)
<b>Counseling for tobacco use</b> ( Ask all adults about tobacco use, advise them to stop using tobacco, and provide behavioral interventions and U.S. Food and Drug Administration–approved pharmacotherapy for cessation to adults who use tobacco AND ask all pregnant women about tobacco use and provide augmented, pregnancy-tailored counseling for those who smoke)	99385-99387 , 99395-99397 or 99406-99407,	Z87.891, Z72.0, Z71.6, Z34.00, Z34.01, Z34.02, Z34.03, Z34.80, Z34.81, Z34.82, Z34.83, Z34.90, Z34.91, Z34.92, Z34.93, Z33.1, O09.00, O09.01, O09.02, O09.03, O09.10, O09.11, O09.12, O09.13, O09.291, O09.292, O09.293, O09.299, O09.40, O09.41, O09.42, O09.43, O09.211, O09.212, O09.213, O09.219, O09.30, O09.31, O09.32, O09.33, O09.511, O09.512, O09.513, O09.519, O09.521, O09.522, O09.523, O09.529, O09.611, O09.612, O09.613, O09.619, O09.621, O09.622, O09.623, O09.629, O09.811, O09.812, O09.813, O09.819, O09.821, O09.822, O09.823, O09.829, O36.80X0, O36.80X1, O36.80X2, O36.80X3, O36.80X4, O36.80X5, O36.80X9, O09.891, O09.892, O09.893, O09.899, O09.70, O09.71, O09.72, O09.73, O09.90, O09.91, O09.92, O09.93, F17.200, F17.201, F17.210, F17.211, F17.220, F17.221, F17.290, F17.291, O99.330, O99.331, O99.332, O99.333, O99.334, O99.335 O09.A0, O09.A1, O09.A2, O09.A3	<a href="#">FDA-approved tobacco cessation prescription medications</a> and OTC nicotine replacement therapy (NRT) are covered at 100%. This is limited to a 90 day supply each for two cessation efforts. NRT is available only with a prescription. Telephonic counseling is available by calling 844-8NCQUIT. <b>NOTE: Z33.1 is not an acceptable principle diagnosis code.</b>
<b>Primary care interventions to prevent tobacco use in children and adolescents</b> (including education or brief counseling to prevent initiation of tobacco use among school-aged children and adolescents)	99383, 99384, 99385, 99393, 99394, 99395, 99401, 99402, 99403, 99404	Z00.121, Z00.129, Z00.110, Z00.111, Z00.00, Z00.01	99401-99404 are to be used in the absence of a wellness visit.
<b>Screening for visual acuity in children</b> (vision screening for all children at least once between the ages of 3 and 5 years to detect the presence of amblyopia or its risk factors)	99382, 99392, 99173, 99174, or 99177	Z01.00, Z01.01, Z13.5	



## Health Care Reform Preventive Services Coding Guide

<b>Bright Futures Recommendations for Children</b>	<b>CPT or HCPCS</b>	<b>ICD-10 Diagnosis</b>	<b>Comments</b>
<b>Sensory Screening - Vision</b>	99381, 99382, 99383, 99384, 99385, 99391, 99392, 99393, 99394, 99395, 99173, 99174 or 99177	Z00.121, Z00.129, Z00.110, Z00.111, Z13.5, Z01.00, Z01.01	May be part of well-child visits.
<b>Sensory Screening - Hearing</b> (beyond newborn screening)	99382, 99383, 99384, 99385, 99392, 99393, 99394, 99395, 92551, 92558 or 92586	Z00.121, Z00.129, Z00.110, Z00.111, Z01.10, Z01.118	May be part of wellness visits.
<b>Developmental Screening</b>	96110	<b>Z13.4</b> Termed 9/30/18 Effective 4/1/18 <b>Z00.121, Z00.129</b> , Effective 10/1/18: <b>Z13.42, Z13.49</b>	96111 is not a screening service and should not be reported for this recommendation.
<b>Autism Screening</b>	96110	<b>Z13.4</b> , Termed 9/30/18 <del><b>Z13.89</b></del> Remove effective 10/1/18 Effective 10/1/18: <b>Z13.41</b>	96111 is not a screening service and should not be reported for this recommendation.
<b>Developmental Surveillance</b>	99381, 99382, 99383, 99384, 99385, 99391, 99392, 99393, 99394, 99395	Z00.121, Z00.129, Z00.110, Z00.111	May be part of well-child visits.
<b>Psychosocial/Behavioral Assessment</b>	99381, 99382, 99383, 99384, 99385, 99391, 99392, 99393, 99394, 99395	Z00.121, Z00.129, Z00.110, Z00.111	Part of well-child visits.
<b>Depression Screening</b> (screening at ages 11-21)	99383, 99384, 99385, 99393, 99394, 99395, G0444	Z00.121, Z00.129, Z00.110, Z00.111, <del>Z13.89</del> Remove effective 9/30/18 Effective 10/1/18: <b>Z13.31</b>	
<b>Alcohol and Drug Use Assessment</b>	99408, 99409, G0442, G0443, G0396 or G0397 Effective 1/1/19: <b>G2011</b>	Z13.89, F10.10, F10.120, F10.129	These codes are to be used when the service is not part of a wellness visit.
<b>Hematocrit or Hemoglobin</b>	85013, 85014, 85018, 85025, 85027	Z00.121, Z00.129, Z00.110, Z00.111, Z13.0	
<b>Congenital Heart Disease Screening</b>	Not applicable, administered through Facility	Not applicable, administered through Facility	Screening for critical congenital heart disease using pulse oximetry should be performed in newborns after 24 hours of age before discharge from the hospital. This screening would be administered to newborn at the birth facility
<b>Lead Screening</b> (up to 7 yrs)	83655	Z13.88	
<b>Tuberculin Test</b>	86580	Z11.1	
<b>Dyslipidemia Screening</b> (Cholesterol)	80061, 82465, or 83718	Z13.220	
<b>STI/HIV Screening</b>	G0432, G0433 or G0435 or G0475	Z34.00, Z34.01, Z34.02, Z34.03, Z34.80, Z34.81, Z34.82, Z34.83, Z34.90, Z34.91, Z34.92, Z34.93, Z33.1, O09.00, O09.01, O09.02, O09.03, O09.10, O09.11, O09.12, O09.13, O09.291, O09.292, O09.293, O09.299, O09.40, O09.41, O09.42, O09.43, O09.211, O09.212, O09.213, O09.219, O09.30, O09.31, O09.32, O09.33, O09.511, O09.512, O09.513, O09.519, O09.521, O09.522, O09.523, O09.529, O09.611, O09.612, O09.613, O09.619, O09.621, O09.622, O09.623, O09.629, O09.811, O09.812, O09.813, O09.819, O09.821, O09.822, O09.823, O09.829, O36.80X0, O36.80X1, O36.80X2, O36.80X3, O36.80X4, O36.80X5, O36.80X9, O09.891, O09.892, O09.893, O09.899, O09.70, O09.71, O09.72, O09.73, O09.90, O09.91, O09.92, O09.93, Z11.4 O09.A0, O09.A1, O09.A2, O09.A3	A screen for HIV has been added between 16 and 18 years. Information on screening adolescents for HIV has been added in the footnotes. STI screening now references recommendations made in the AAP Red Book. This category was previously titled, "STI Screening". <b>NOTE:</b> Z33.1 is not an acceptable principle diagnosis code.
<b>Oral Health</b>	99381, 99382, 99383, 99384, 99391, 99392, 99393, 99394	Z00.121, Z00.129, Z00.110, Z00.111	May be part of well-child visits.
<b>Anticipatory Guidance</b>	99381, 99382, 99383, 99384, 99391, 99392, 99393, 99394	Z00.121, Z00.129, Z00.110, Z00.111	May be part of well-child visits.
<b>Bilirubin Screening</b> Universal assessment of bilirubin level in infants with gestational age of 35 weeks or greater	82247, 88720	Z00.110	Mandate effective 1/1/18 or upon renewal Service is typically performed in the birth facility or as part of a wellness office visit in the event of a home birth.
<b>Postpartum depression screening</b> Performed as part of 1, 2, 4, and 6 month well baby visits	96161	Z00.110, Z00.111, Z00.121, Z00.129 Effective 10/1/18: <b>Z13.32</b>	Mandate effective 1/1/18 or upon renewal To be billed as part of newborn/infant visit. Not mother's postpartum visit



## Health Care Reform Preventive Services Coding Guide

<b>Women's Preventive Services effective at renewal on or after August 1, 2012, unless otherwise noted</b>	<b>CPT or HCPCS</b>	<b>ICD-10 Diagnosis</b>	<b>Frequency</b>
Well Woman visits (Well-woman preventive care visit annually for adult women to obtain the recommended preventive services that are age and developmentally appropriate, including preconception and prenatal care. This well-woman visit should, where appropriate, include other preventive services listed in this set of guidelines.	99384, 99385, 99386, 99387 or 99394, 99395, 99396, 99397	Z00.00, Z00.01, Z01.411, Z01.419	Annual, although HHS recognizes that several visits may be needed to obtain all necessary recommended preventive services, depending on a woman's health status, health needs, and other risk factors
Screening for gestational diabetes	82950	Z34.00, Z34.01, Z34.02, Z34.03, Z34.80, Z34.81, Z34.82, Z34.83, Z34.90, Z34.91, Z34.92, Z34.93, Z33.1, O09.00, O09.01, O09.02, O09.03, O09.10, O09.11, O09.12, O09.13, O09.291, O09.292, O09.293, O09.299, O09.40, O09.41, O09.42, O09.43, O09.211, O09.212, O09.213, O09.219, O09.30, O09.31, O09.32, O09.33, O09.511, O09.512, O09.513, O09.519, O09.521 O09.522, O09.523, O09.529, O09.611, O09.612, O09.613, O09.619, O09.621, O09.622, O09.623, O09.629, O09.811, O09.812, O09.813, O09.819, O09.821, O09.822, O09.823, O09.829, O36.80X0, O36.80X1, O36.80X2, O36.80X3, O36.80X4, O36.80X5, O36.80X9, O09.891, O09.892, O09.893, O09.899, O09.70, O09.71, O09.72, O09.73, O09.90, O09.91, O09.92, O09.A0, O09.A1, O09.A2, O09.A3	In pregnant women between 24 and 28 weeks of gestation and at the first prenatal visit for pregnant women identified to be at high risk for diabetes. <b>NOTE: Z33.1 is not an acceptable principle diagnosis code.</b>
Human papillomavirus (HPV) testing (High-risk human papillomavirus DNA testing in women with normal cytology results.)	87623, 87624 , 87625 or G0476	Z11.51	Screening should begin at 30 years of age and should occur no more frequently than every 3 years.
Counseling for sexually transmitted infections for all sexually active women	Part of preventive visit, 99401-99404 or G0445	Z71.7, Z71.89, Z72.51, Z72.52, Z72.53	Annual. 99401-99404 are to be used in the absence of a wellness visit.
Counseling and screening for human immunodeficiency virus for all sexually active women	<u>Counseling:</u> part of preventive visit or 99401, 99402, 99403, 99404 <u>Screening:</u> G0432, G0433 or G0435 or G0475	Z34.00, Z34.01, Z34.02, Z34.03, Z34.80, Z34.81, Z34.82, Z34.83, Z34.90, Z34.91, Z34.92, Z34.93, Z33.1, O09.00, O09.01, O09.02, O09.03, O09.10, O09.11, O09.12, O09.13, O09.291, O09.292, O09.293, O09.299, O09.40, O09.41, O09.42, O09.43, O09.211, O09.212, O09.213, O09.219, O09.30, O09.31, O09.32, O09.33, O09.511, O09.512, O09.513, O09.519, O09.521 O09.522, O09.523, O09.529, O09.611, O09.612, O09.613, O09.619, O09.621, O09.622, O09.623, O09.629, O09.811, O09.812, O09.813, O09.819, O09.821, O09.822, O09.823, O09.829, O36.80X0, O36.80X1, O36.80X2, O36.80X3, O36.80X4, O36.80X5, O36.80X9, O09.891, O09.892, O09.893, O09.899, O09.70, O09.71, O09.72, O09.73, O09.90, O09.91, O09.92, O09.93, Z71.7, Z11.4 O09.A0, O09.A1, O09.A2, O09.A3	Annual. 99401-99404 are to be used in the absence of a wellness visit. <b>NOTE: Z33.1 is not an acceptable principle diagnosis code.</b>
Contraceptive method ( including-but not limited to, sterilization and IUD), and counseling (Group health plans sponsored by certain religious employers, and group health insurance coverage in connection with such plans, are exempt from the requirement to cover contraceptive services)	Part of preventive visit (99384, 99385, 99386, 99394, 99395, 99396), 99401, 99402, 99403, 99404, S0610, S0612, S0613, A4261, A4264, J7300, J7302, J7306, J7307, J1050, S4981, S4989, 11981, 11982, 11983, 57170, 58300, 58565, 58600, 58605, 58611, 58615, 58671, 58670, 96372, 11976, 58301, 74740, 58340, J7301, 00952, 00851, J7297, J7298 or -J7296	Z34.00, Z34.01, Z34.02, Z34.03, Z34.80, Z34.81, Z34.82, Z34.83, Z34.90, Z34.91, Z34.92, Z34.93, Z33.1, O09.00, O09.01, O09.02, O09.03, O09.10, O09.11, O09.12, O09.13, O09.291, O09.292, O09.293, O09.299, O09.40, O09.41, O09.42, O09.43, O09.211, O09.212, O09.213, O09.219, O09.30, O09.31, O09.32, O09.33, O09.511, O09.512, O09.513, O09.519, O09.521 O09.522, O09.523, O09.529, O09.611, O09.612, O09.613, O09.619, O09.621, O09.622, O09.623, O09.629, O09.811, O09.812, O09.813, O09.819, O09.821, O09.822, O09.823, O09.829, O36.80X0, O36.80X1, O36.80X2, O36.80X3, O36.80X4, O36.80X5, O36.80X9, O09.891, O09.892, O09.893, O09.899, O09.70, O09.71, O09.72, O09.73, O09.90, O09.91, O09.92, O09.93, Z30.011, Z30.013, Z30.014, Z30.017, Z30.018, Z30.019, Z30.012, Z30.02, Z30.09, Z30.13, Z30.430, Z30.433, Z30.2, Z30.40, Z30.41, Z30.431, Z30.49, Z30.42, Z30.46, Z30.49, Z30.8, Z30.9, Z30.432 O09.A0, O09.A1, O09.A2, O09.A3	Diaphragms, vaginal rings, contraceptive patches, female condoms, sponges, spermicides, and emergency contraception are available at a participating pharmacy with an MD prescription.  Anesthesia services for 00952 and 00851 will be provided with no cost sharing to appropriate groups If there is a medical reason a member cannot take a generic contraceptive, the physician should review the clinical <a href="#">criteria</a> and if appropriate, submit fax <a href="#">form</a> .  99401-99404 are to be used in the absence of a wellness visit. Injectable contraceptives and administration (J1050, 96372) are covered when billed with  <b>NOTE: Z33.1 is not an acceptable principle diagnosis code.</b>

## Health Care Reform Preventive Services Coding Guide

<b>Women's Preventive Services effective at renewal on or after August 1, 2012, unless otherwise noted</b>	<b>CPT or HCPCS</b>	<b>ICD-10 Diagnosis</b>	<b>Frequency</b>
Breastfeeding support, supplies, and counseling (Comprehensive lactation support and counseling, by a trained provider during pregnancy and/or in the postpartum period, and costs for breastfeeding equipment)	99401, 99402, 99403, S9443, E0602, E0603, A4281, A4282, A4283, A4284, A4285, A4286	O30.93, Z34.03, Z34.83, Z34.93, Z39.1	Counseling covered at 100% through in-network providers (i.e., OB/GYNs, midwives, facilities); one breast pump provided per pregnancy through in-network DME providers. Replacement supplies for breast pumps or initial supplies for member owned breast pump will be provided with no cost sharing to appropriate groups <b>Replacement supplies cannot be provided with the initial breast pump as they are included in the initial purchase of the pump</b> 99401-99404 are to be used in the absence of a wellness visit.
Screening and counseling for interpersonal and domestic violence	Part of preventive visit or 99401, 99402, 99403, 99404	Z00.00, Z00.01, Z01.411, Z01.419	Annual 99401-99404 are to be used in the absence of a wellness visit.
<b>Effective 1/1/19 or upon renewal</b> Screening for diabetes after pregnancy (For women with a history of gestational diabetes who are not currently pregnant and who have not previously been diagnosed with type 2 diabetes. Frequency: Initial testing 3 should ideally occur within the first year postpartum and can be conducted as early as 4 to 6 weeks postpartum)	82947, 83036 or 82950	:Z86.32	
<b>Effective 1/1/19 or upon renewal</b> Screening for Urinary Incontinence (Screening should assess whether women experience urinary incontinence and whether it impacts their activities and quality of life. Clinicians should refer women for further evaluation and treatment if indicated )	Part of preventive visit or 99401, 99402, 99403, 99404	Z00.00, Z00.01, Z01.411, Z01.419	Annual 99401-99404 are to be used in the absence of a wellness visit.

<b>Vaccines Recommended by the Centers for Disease Control (CDC)</b>	<b>ICD-10 Diagnosis</b>	<b>Comments</b>
<p>Diphtheria, Tetanus, Pertussis Measles, Mumps, Rubella Haemophilus Infuenzae Type B Hepatitis A Hepatitis B Influenza Pneumococcal Meningococcal Human Papillomavirus (HPV)*</p> <p>Inactivated Poliovirus Rotavirus Varicella Tetanus-Diphtheria /Tetanus-Diphtheria Acellular Pertussis Herpes Zoster (Shingles)</p> <p>Adult and Child &amp; Adolescent Immunization Schedules (for persons aged 0-6 years, 7-18 years, and "catch-up schedule")</p> <p>Refer to the CDC's posted schedule of immunizations <a href="http://www.cdc.gov/vaccines/schedules/index.html">http://www.cdc.gov/vaccines/schedules/index.html</a></p> <p>*The HPV vaccine (Gardasil) is covered to age 45 per the Food and Drug Administration (FDA) guidelines</p>	Z23	Doses, recommended ages and recommended populations vary. All recommended routine immunizations will be allowed with no cost share.

## Health Care Reform Preventive Services Coding Guide

### Revision History January 1, 2017 to July 1, 2019

Date of change	Mandate	Change
January 1, 2017	Screening for Abdominal Aortic Aneurysm	Addition of 76706 effective 1/1/17
	Hypertension in Adults-screening and home monitoring	Added coverage of home monitoring (ABPM). Add DX R03.0, CPT 93784,93786, 93788, 93790 and HCPCS A4670 for ABPM with criteria
	Screening for Breast Cancer	Added 77067 effective 1/1/18. Removed 77052, 77057 termed 12/31/16
	Colorectal Cancer Screening	Added new HCPCS code G0500 with following instructions: Moderate sedation (G0500) performed by the physician performing the procedure is allowed but cannot be billed on the same DOS as 00810
June 1, 2017	Screening for Breast Cancer	Addition of coverage for 3-D tomosynthesis and CPT code 77063
July 1, 2017	Breast Feeding Support, Supplies and Counseling	Added statement clarifying that replacement supplies cannot be provided at same time as initial breast pump as they are included in purchase of pump
October 1, 2017		Remove ICD-9 codes from all mandates
	<b>Screening for latent tuberculosis</b>	<b>New mandate effective 10/1/17</b>
	Counseling related to BRCA screening	Addition of diagnosis code Z71.83 effective 10/1/17. Removed Z31.5 as no longer applicable
	Screening for chlamydial infection in non-pregnant and pregnant women	Removed 87800 and 87801 as more specific codes should be used to report service
	Screening for gonorrhea for women	Removed 87800 and 87801 as more specific codes should be used to report service
	Vaccines	Added Shingrix to covered vaccines
December 1, 2017	Statin Use for the Primary Prevention of Cardiovascular Disease in Adults	New mandate effective 12/1/17
January 1, 2018	<b>Bilirubin Screening</b>	<b>New mandate effective 1/1/18 or upon group renewal</b>
	<b>Postpartum depression screening</b>	<b>New mandate effective 1/1/18 or upon group renewal</b>
	Colorectal cancer screening	Added 00812 effective 1/1/18. Removed 00810 termed 12/31/17
	Contraceptives	Added J7296 effective 1/1/18. Removed Q9984 termed 12/31/17
	Screening for Breast Cancer	Removed G0202 termed 12/31/17
	Screening for Cervical Cancer	Removed 88154 termed 12/31/17
April 1, 2018	Developmental screening	Added dx codes Z00.129 and Z00.120 to covered diagnosis codes effective 4/1/18
May 1, 2018	<b>Screening for preeclampsia</b>	<b>New mandate effective 5/1/18 or upon group renewal</b>
October 1, 2018	Postpartum depression screening	Added dx code Z13.32 effective 10/1/18
	Depression screening (adult and Bright Futures)	Added dx code Z13.31 effective 10/1/18. Removed Z13.89 effective 9/30/18 (should no longer be used as more specific code available)
	Autism screening	Added dx code Z13.41 effective 10/1/18. Removed Z13.89 effective 9/30/18 (should no longer be used as more specific code available)
January 1, 2019	<b>Screening for Diabetes after Pregnancy</b>	<b>Added new mandated coverage for screening in women with previous diagnosis of gestational diabetes effective 1/1/19 or upon group renewal</b>
	Screening and counseling to reduce alcohol misuse	Added HCPCS code G2011 effective 1/1/19
	Counseling related to BRCA screening	Added CPT codes 81163, 81164, 81165, 81166, 81167 effective 1/1/19. Removed CPT codes 81211, 81213, 81214 effective 12/31/18.
	<b>Screening for Urinary Incontinence</b>	<b>New mandate effective 1/1/19 or upon group renewal</b>
March 1, 2019	Contraceptives	Added Z30.017 and Z30.46
April 1, 2019	Fall Prevention	Revised description based on updated USPSTF recommendation
	Behavioral Counseling to Prevent Skin Cancer	Revised description based on updated USPSTF recommendation
July 1 2019	Screening for Osteoporosis	Revised description based on updated USPSTF recommendation
August 1, 2019	Colorectal Cancer Screening	Added 81528 (Cologuard) to covered tests.